

Separate Drop Schedule

For each eye you will be prescribed 3 drops to your pharmacy. Drops are **not** available at the surgery center. To keep your eyes moist, we recommend lubricating drops. Coupons are available at all Eyesight locations.



Morning of Surgery: Starting 1 hour before leaving home, use 1 drop each of the moxifloxacin, prednisolone, and ketorolac every 15 minutes for a total of 4 doses. i.e. if you will be leaving home at 9:00 AM, use the drops at 8:00, 8:15, 8:30, and 8:45 AM. Shake bottles before use. **Separate drops by at least 5 minutes.** Do not share bottles between eyes.

Beginning 1 HOUR BEFORE LEAVING HOME FOR THE SURGERY CENTER on _____

1 hour prior	45 minutes prior	30 minutes prior	15 minutes prior	After leaving the surgery center, use again at 12:00pm – 4:00pm – 8:00pm
□	□	□	□	

After Surgery: (4 times a day is roughly 8am, 12pm, 4pm, 8pm. 2 times a day is roughly 8am and 8pm)

Week 1	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
Moxifloxacin or Polytrim	4 times a day □□□□	4 times a day □□□□	4 times a day □□□□	4 times a day □□□□	4 times a day □□□□	4 times a day □□□□	4 times a day □□□□
Prednisolone Acetate 1%	4 times a day □□□□	4 times a day □□□□	4 times a day □□□□	4 times a day □□□□	4 times a day □□□□	4 times a day □□□□	4 times a day □□□□
Ketorolac	4 times a day □□□□	4 times a day □□□□	4 times a day □□□□	4 times a day □□□□	4 times a day □□□□	4 times a day □□□□	4 times a day □□□□

Week 2	Day 8	Day 9	Day 10	Day 11	Day 12	Day 13	Day 14
Prednisolone Acetate 1%	4 times a day □□□□	4 times a day □□□□	4 times a day □□□□	4 times a day □□□□	4 times a day □□□□	4 times a day □□□□	4 times a day □□□□
Ketorolac	4 times a day □□□□	4 times a day □□□□	4 times a day □□□□	4 times a day □□□□	4 times a day □□□□	4 times a day □□□□	4 times a day □□□□
Artificial Tears	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□

Week 3	Day 15	Day 16	Day 17	Day 18	Day 19	Day 20	Day 21
Prednisolone Acetate 1%	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□
Ketorolac	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□
Artificial Tears	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□

Week 4	Day 22	Day 23	Day 24	Day 25	Day 26	Day 27	Day 28
Prednisolone Acetate 1%	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□
Ketorolac	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□
Artificial Tears	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□

It is EXTREMELY important to follow your eyedrop instructions!
NOTE: Your surgeon will discuss the recommended continuation of your eyedrops after 4 weeks

Week 5 & Week 6	Day 29	Day 30	Day 31	Day 32	Day 33	Day 34	Day 35
Prednisolone Acetate 1%	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□
Ketorolac	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□
Artificial Tears	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□	2 times a day □□